

Integer quantum Hall effect and Hofstadter's butterfly spectra in three-dimensional metals in external periodic modulations

Mikito Koshino*, Hideo Aoki

Department of Physics, University of Tokyo, Hongo, Tokyo 113-0033, Japan
(February 2, 2008)

We propose that Hofstadter's butterfly accompanied by quantum Hall effect that is similar to those predicted to occur in 3D tight-binding systems by Koshino *et al.* [Phys. Rev. Lett. **86**, 1062 (2001)] can be realized in an entirely different system — 3D metals applied with weak external periodic modulations (e.g., acoustic waves). Namely, an effect of two periodic potentials interferes with Landau's quantization due to an applied magnetic field $\mathbf{B}$, resulting generally in fractal energy gaps as a function of the tilting angle of $\mathbf{B}$, for which the accompanying quantized Hall tensors are computed. The phenomenon arises from the fact that, while the present system has a different physical origin for the butterfly from the 3D tight-binding systems, the mathematical forms are remarkably equivalent.

Introduction In two-dimensional(2D) periodic systems in magnetic fields, it is well-known that the interplay of Bragg's reflection and Landau's quantization gives rise to a fractal series of energy gaps, which is called Hofstadter's butterfly¹. Theoretically, the self-similar spectrum emerges both in the strong potential limit¹ (tight-binding system) and in the weak potential limit² (Landau-quantized system), where two cases relate to each other in their mathematical expressions. When the Fermi energy is within a gap in the butterfly the system exhibits the quantum Hall effect, where the Hall conductivity is expressed as a topological quantum number for each energy gap³. In recent experiments, a signature of Hofstadter's butterfly has been reported in the Hall conductivity measurement for semiconductor superlattices⁴.

While the butterfly spectra had been known to be peculiar to 2D, we have previously found that *three-dimensional(3D) tight-binding lattice* can have a fractal energy spectra like Hofstadter's when the system is anisotropic (quasi-1D), where the 3D butterfly is shown to be a genuinely 3D effect rather than a remnant of a 2D butterfly⁵. This leads to an interesting transport phenomenon — quantum Hall effect (QHE) in three dimensions. According to the general argument^{6–8}, QHE occurs even in 3D *provided* that there is an energy gap and that the Fermi energy lies in the gap, where the three components of the Hall conductivity tensor should be individually quantized.

Then an intriguing question arises: can we have a counterpart in the opposite limit, *3D weakly modulated systems*? Here we investigate the energy spectra in 3D continuous systems with weak periodic potentials, and show that a modulation composed of two plane waves (such as two acoustic waves) gives rise to Hofstadter's butterfly, although the mechanism is quite distinct from one in the tight-binding system. We also find a clear mathematical relationship connecting two limiting cases similarly to 2D, but we will see that three-dimensionality is essential throughout.

The problem has another interesting aspect, i.e., the QHE topological numbers. Here we have calculated the

Hall conductivity in 3D weak potential systems, and find that the expression is remarkably equivalent to that for the corresponding energy gap in the tight-binding system⁵, which is unlike the relationship in 2D between the limits of weak $\leftrightarrow$ strong potentials.

The present experimental setup (just an ordinary 3D metal with two acoustic waves applied) is so simple that this can be a strong candidate for experimentally detecting the 3D QHE and Hofstadter's butterfly. We shall estimate the required magnetic field and acoustic wave length, etc, for a possible realization in a semimetal subjected.

Formulation: One modulation We first look at how a single modulation interferes with Landau's quantization due to a magnetic field $\mathbf{B}$ in a 3D uniform electron gas. The Hamiltonian is simply $\mathcal{H} = (1/2m)(\mathbf{p} + e\mathbf{A})^2 + U(\mathbf{r})$, where $U(\mathbf{r})$ is a perturbative periodic potential, and the vector potential is taken as $\mathbf{A} = (0, Bx, 0)$ for $\mathbf{B} = (0, 0, B)$. The eigenstates in the absence of $U(\mathbf{r})$ are, as usual,

$$|n, k_y, k_z\rangle = N_n \exp(ik_y y + ik_z z) \Psi_n(x/l + k_y l),$$

$$\Psi_n(z) = e^{-z^2/2} H_n(z), \quad (1)$$

where H_n is the Hermite polynomial with the Landau index n , N_n a normalization factor, and $l = \sqrt{\hbar/(eB)}$ is the magnetic length (~ 80 Å for $B = 10$ T). The eigenenergy is $E_{n, k_z} = (n + 1/2)\hbar\omega_c + \hbar^2 k_z^2/(2m)$ with the electron mass m and the cyclotron frequency $\omega_c = eB/m$, which suggests that a 3D uniform electron gas becomes 1D in that the motion along the magnetic field remains free. So, if we have an external modulation on top, we immediately expect energy gaps to emerge in the 1D band.

When we apply a single periodic potential $U(\mathbf{r}) = U_0 \cos(\mathbf{G} \cdot \mathbf{r})$ in 3D, a mixing between the states $|n, k_y, k_z\rangle$ and $|n', k_y \pm MG_y, k_z \pm MG_z\rangle$ gives rise to energy gaps in the M -th order perturbation. The energy gaps due to the mixing between higher Landau bands will generally tend to be hidden by other bands, so we focus on the lowest two levels. Namely, energy gaps should open within the lowest Landau level ($n = n' = 0$) and between

the lowest and second levels ($n = 0, n' = 1$) (see, Fig. 1). The corresponding matrix elements in the first order perturbation are

$$\begin{aligned} \langle 0, k_y \pm G_y, k_z \pm G_z | U(\mathbf{r}) | 0, k_y, k_z \rangle \\ = U_0 e^{-G_\perp^2 l^2 / 4} e^{\mp i(k_y \pm \frac{G_y}{2}) G_x l^2}, \end{aligned} \quad (2)$$

$$\begin{aligned} \langle 1, k_y \pm G_y, k_z \pm G_z | U(\mathbf{r}) | 0, k_y, k_z \rangle \\ = U_0 e^{-G_\perp^2 l^2 / 4} e^{\mp i(k_y \pm \frac{G_y}{2}) G_x l^2} \times i(G_x - iG_y)l/\sqrt{2} \end{aligned} \quad (3)$$

where $G_\perp \equiv \sqrt{G_x^2 + G_y^2}$. Since the matrix elements scale as $\exp(-G_\perp^2 l^2 / 4)$, the magnitude of the matrix elements, hence the size of the energy gap, becomes significant when $Gl \lesssim 1$ (i.e., $\exp(-G_\perp^2 l^2 / 4)$ vanishes like $\exp(-\text{const.}/B)$ for $B \rightarrow 0$). The condition $Gl \lesssim 1$ also suggests that the main energy gaps open for $E \lesssim \hbar\omega_c$, i.e., around the lowest or the second lowest Landau levels.

In 3D systems, the Hall conductivity has three components, $\boldsymbol{\sigma} \equiv (\sigma_{yz}, \sigma_{zx}, \sigma_{xy})$, where the Hall current in an electric field $\mathbf{E}$ is given by $\mathbf{j} = \boldsymbol{\sigma} \times \mathbf{E}$. We can immediately calculate $\boldsymbol{\sigma}$ for the present system when the Fermi energy E_F is in each of the energy gaps. If we consider generally the M -th order gap ($M = 1, 2, \dots$) within the lowest Landau level ($n = n' = 0$), the number of states below E_F is

$$N_F = \frac{MG_z}{2\pi/L_z} \times \frac{L_x L_y}{2\pi l^2} = \frac{eV}{2\pi\hbar} M \mathbf{G} \cdot \mathbf{B}, \quad (4)$$

where V is the system volume. From Widom-Středa's formula^{10,11}, $\boldsymbol{\sigma} = -(e/V)(\partial N_F / \partial \mathbf{B})$, we readily obtain

$$\boldsymbol{\sigma} = -\frac{e^2}{2\pi\hbar} M \mathbf{G} \quad (n = n' = 0). \quad (5)$$

Similarly, we have for the inter-Landau gap

$$\boldsymbol{\sigma} = -\frac{e^2}{2\pi\hbar} 2M \mathbf{G} \quad (n = 0, n' = 1). \quad (6)$$

A note is due here. 3D continuous systems having a one-dimensional modulation in magnetic fields have been investigated for a long time in terms of density-wave (DW) instabilities. Specifically, Halperin⁶ has calculated the Hall conductivity when the system is in a DW state, which corresponds to our calculation for the energy gap within the lowest Landau level. The *inter-Landau level* gap formation is new to the best of our knowledge.

Two modulations Now, if we superpose two modulations $U_1(\mathbf{r}) = U_1 \cos(\mathbf{G}_1 \cdot \mathbf{r})$ and $U_2(\mathbf{r}) = U_2 \cos(\mathbf{G}_2 \cdot \mathbf{r})$ having different wavevectors, the discussion above can be extended, where we can substitute $M\mathbf{G}$ with $M\mathbf{G}_1 + N\mathbf{G}_2$ for the energy gap corresponding to M th and N th-order perturbations in U_1 and U_2 , respectively. The Hall conductivity for the gap with the order M, N in the lowest Landau band thus becomes

$$\boldsymbol{\sigma} = -\frac{e^2}{2\pi\hbar} (M\mathbf{G}_1 + N\mathbf{G}_2) \quad (n = 0, n' = 0), \quad (7)$$

where (M, N) becomes $(2M, 2N)$ for the (M, N) -th inter-Landau level gap.

We can immediately realize that the spectrum should be sensitively affected by the commensurability of the ratio between the z -components, G_{1z} and G_{2z} , since, for instance if the ratio is irrational $MG_{1z} + NG_{2z}$ can take continuous values and the spectrum should have gaps everywhere in the energy axis. If the magnetic field is *rotated* in the plane spanned by $\mathbf{G}_1$ and $\mathbf{G}_2$, the energy spectrum is then expected to have a fractal structure since the commensurability between G_{1z} and G_{2z} varies in a complicated manner.

The spectrum plotted in Fig.2 against the tilting angle of $\mathbf{B}$ obtained numerically, has indeed butterfly-like structure in the lowest Landau band as in Hofstadter's spectrum. The Hall integers M, N have been calculated rigorously for each gap, by tracing the number of states below the gaps for the tilted field. In higher energy region where two Landau levels overlap ($E > (3/2)\hbar\omega_c$) we see the gap structure due to the inter-Landau level mixing (the gap labeled with $M, N = 4, 0(0, 4)$ and $2, 2$ in the figure), as well as the intra-level gaps which are visible only when gaps within $n = 0$ and $n = 1$ happen to overlap. In the figure, the gap $3, 1(1, 3)$ is a composite of $2, 1(1, 2)$ in the lowest level and $1, 0(0, 1)$ in the second.

Correspondence with the strong potential case As mentioned, we have previously found a butterfly-like spectra for the first time in periodic 3D systems which is modelled by the anisotropic (quasi-1D) *tight-binding* lattice. There we have shown that the period along the most conductive direction (x) is *not* responsible for the emergence of the 3D butterfly, while other two periods (along y, z) are relevant. Thus the two (present and the previous) cases are both *3D system with two periods* and the only difference is the strength of the periodic potential. One might then be tempted to think that the two butterfly spectra cross over to each other when the amplitude of the periodic modulation is increased or decreased. However, as shown in the following, they have distinct physical origins, residing in different limits, and the resemblance in the spectra comes from a mathematical relationship.

To show this, let us concentrate on weak modulations with $G_1, G_2 \lesssim 1/l$ for the lowest Landau level (i.e., low-energy region of the spectrum). To clarify the relationship, we concentrate on a case where $\mathbf{B}$ lies on the plane spanned by $\mathbf{G}_1, \mathbf{G}_2$. If we take $\mathbf{G}_1$ and $\mathbf{G}_2$ on zx -plane (with $G_{1y} = G_{2y} = 0$) so that k_y is conserved, Schrödinger's equation becomes

$$\frac{\hbar^2 k_z^2}{2m} \psi(k_z) + \sum_{k'_z} U_{k_z, k'_z} \psi(k'_z) = E \psi(k_z), \quad (8)$$

where $\psi(k_z)$ is the amplitude of $|0, k_y, k_z\rangle$ with k_y being constant, U_{k_z, k'_z} is eq.(2) summed over U_1 and U_2 with $G_y = 0$, and is a function of k_y . By Fourier-transforming with respect to z we obtain

$$-\frac{\hbar^2}{2m} \frac{\partial^2}{\partial z^2} \psi(z) + \tilde{U}_1 \cos(G_{1z}z - k_y G_{1x}l^2) \psi(z) + \tilde{U}_2 \cos(G_{2z}z - k_y G_{2x}l^2) \psi(z) = E\psi(z), \quad (9)$$

where $\tilde{U}_j = U_j e^{-G_{j\perp}^2 l^2/4}$ with $G_{j\perp} \equiv \sqrt{G_{jx}^2 + G_{jy}^2}$ ($j = 1, 2$).

Now let us recapitulate how a butterfly spectrum arose in our original model on a 3D tight-binding lattice⁵. We consider the 3D orthorhombic lattice with the nearest-neighbor transfers t_x, t_y, t_z along x, y, z , respectively. We assume here that the magnetic field $\mathbf{B}$ is applied parallel to yz -plane ($B_x = 0$), which corresponds to the assumption made in the weak potential case that $\mathbf{G}_1, \mathbf{G}_2, \mathbf{B}$ are co-planar. We also assume that the system is quasi-1D ($t_x \gg t_y, t_z$), and apply the effective mass approximation to the motion along the conductive direction x , to obtain the low-energy spectra ($\lesssim t_y$ or t_z from the bottom). In a gauge $\mathbf{A} = (0, B_z x, -B_y x)$ we can take the basis $\Psi(\mathbf{r}) = e^{ik_y y + ik_z z} \psi(x)$, and Schrödinger's equation becomes one-dimensional^{7,5}:

$$-\frac{\hbar^2}{2m} \frac{\partial^2}{\partial x^2} \psi(x) - 2t_y \cos\left(\frac{eB_z b}{\hbar} x + k_y b\right) \psi(x) - 2t_z \cos\left(-\frac{eB_y c}{\hbar} x + k_z c\right) \psi(x) = E\psi(x), \quad (10)$$

where b, c are the lattice constants along y, z , respectively. Now we can see that eq.(9) and (10) have the identical form: 1D equation with a double period. We should recall that Hofstadter's butterfly emerges generally in the spectrum of a doubly periodic 1D system when the spectrum is plotted against the ratio of the two periods⁵. In each of two equations, the ratio changes continuously when the magnetic field is rotated relative to the periodic potential, which is why Hofstadter's butterfly arises against the tilting angle in both cases.

From the correspondence we can also note that $\tilde{U}_1, \tilde{U}_2$ in eq.(9) correspond to t_y, t_z in (10), respectively, but their physical meaning is opposite in the following sense. Namely, $\tilde{U}_1, \tilde{U}_2 \rightarrow 0$ describes the limit of weak periodic potential, while $t_y, t_z \rightarrow 0$ the strong limit. The equations become purely 1D in both limits, where the system reduces to 'Landau tubes' (1D motion along $\mathbf{B}$ and the cyclotron motion on the plane $\perp \mathbf{B}$) in the former, while in the latter wires along x confined by the strong potential on yz plane. On the other hand, if the perturbations ($\tilde{U}_1, \tilde{U}_2$ or t_y, t_z) are too large, the mixing between different 1D channels (i.e., different Landau levels in the former, different bound modes on yz plane in the latter) becomes so strong that the one-band approximation breaks down. We thus conclude that the two butterfly spectra in fact correspond to opposite limits (weak/strong potential).

We can also establish a relationship between the Hall conductivities between the two cases. In the strong potential the Hall conductivity for the gap with M -th in t_y and the N -th in t_z in the tight-binding band can be obtained similarly as⁵

$$\sigma = -\frac{e^2}{h} \left(0, \frac{N}{b}, \frac{M}{c}\right). \quad (11)$$

Since the primitive reciprocal lattice vectors are $\mathbf{G}_1 = (2\pi/c)\hat{e}_z$ and $\mathbf{G}_2 = (2\pi/b)\hat{e}_y$, this coincides with the corresponding expression in the weak potential, eq.(7). This is rather remarkable, since the corresponding wave functions in the two cases have totally different spatial behaviors, while they carry the identical Hall current. This contrasts with the 2D case, where the Hall integers are different between the corresponding gaps in the weak- and strong-potential limits.




FIG. 1. The energy dispersion plotted against the wavenumber along z ($\parallel \mathbf{B}$) in a continuous 3D metal in a magnetic field $\mathbf{B}$ applied in a general direction (inset). Dashed arrows indicate where the modulation having G_z (the z -component) makes the dispersion gapped.

Some notes on the quantum Hall effect are due here. First, the non-monotonic behavior in the Hall conductivity as E_F is increased (see, Fig. 2) should be a hallmark of the 3D butterfly spectrum as in the 2D butterfly. As for the quantum Hall plateaus, we have to consider the localization effect in the presence of disorder. While this is an interesting future problem, we speculate that each subband would evolve into the localized and the extended states with mobility edges since we have a 3D system here, and that the Hall conductivity would be constant as long as E_F stays in the localized region. For the usual 2D butterfly, a numerical study for a dirty, finite system¹² shows that we still have a nonmonotonic behavior as a sign for the butterfly when the disorder is not too strong, so we expect a similar behavior in the disordered 3D butterfly as well.

Second, while our calculation of the Hall conductivity is based on the bulk description, a finite system has edge states in the bulk gaps, which also contribute to the Hall current. We have previously shown¹³ that the Hall conductance in a finite 3D system is still quantized, in the 3D QHE condition, when we take into account both of the contributions from the surface current (which we have called the wrapping current) and from the bulk current, where the quantized values exactly coincides with those for the infinite system. The argument is quite general, and applies to the present problem as well.




FIG. 2. The energy spectra for 3D continuous systems in two periodic modulations having wavevectors $\mathbf{G}_1, \mathbf{G}_2$ (top), along with the plot for one modulation $\mathbf{G}_1$ (bottom), for $|\mathbf{G}_i| = \sqrt{2}/l$, $\mathbf{G}_1 \perp \mathbf{G}_2$, and the amplitudes of the periodic potentials of $0.2\hbar\omega_c$. The horizontal axis is the angle of $\mathbf{B}$ rotated in the plane spanned by $\mathbf{G}_1$ and $\mathbf{G}_2$. A pair of integers (M, N) attached to each energy gap represents the Hall integers, eq.(7). The spectrum has been plotted for rational angles, $\tan \theta = p/q$ (p, q : mutually prime integers), with an energy cut-off at $4.5\hbar\omega_c$. The arrows on the right indicate the energy regions for the Landau bands in the un-modulated case.

Experimental feasibility Finally let us comment on the experimental feasibility for the weakly modulated QHE system proposed here. To observe the QHE, E_F have to reside in an energy gap, which appears only in the lowest Landau levels. So we require the situation where only a few levels are occupied in 3D (the quantum limit), for which we need a large magnetic field or a small electron concentration. We expect that semimetals should be suitable for the latter condition. We can estimate the required magnetic field for the quantum limit in bismuth, for example, to be $B \gtrsim 10\text{T}$, which is quite modest. More stringent is the condition for the modulation, since the wave length should be such that the Fermi level is in the gap created by the modulation. If we take the acoustic wave for the periodic potential and plug in the velocity of sound for Bi, we have $f \sim 100\text{GHz}$ for $B = 10\text{T}$, which is rather high, although the required frequency ($\propto$ Fermi wavenumber) decreases with B like $f \propto 1/B$. The coupling of the electrons with the acoustic wave, on the other hand, is dominated by the piezoelectric coefficient of the material, so compounds may be advantageous in this respect. An obvious advantage of the externally modulated system proposed here over the tight-binding system is that we can change the wavelength of the external modulation at our disposal. So we expect that we can find a wider possibility for observing Hofstadter's butterfly with the 3D QHE.

* Present address: Department of Physics, Tokyo Institute of Technology, 2-12-1 Ookayama, Meguro, Tokyo 152-8551, Japan.

- ¹ D. R. Hofstadter, Phys. Rev. B **14**, 2239 (1976).
- ² A. Rauh, Phys. Status Solidi B **69**, K9 (1975).
- ³ D. J. Thouless, M. Kohmoto, P. Nightingale, and M. den Nijs, Phys. Rev. Lett **49**, 405 (1982).
- ⁴ C. Albrecht, J.H.Smet, K. v. Klitzing, D. Weiss, V. Umansky and H. Schweizer, Phys. Rev. Lett. **86**, 147 (2001).
- ⁵ M. Koshino, H. Aoki, K. Kuroki, S. Kagoshima, and T. Osada, Phys. Rev. Lett. **86**, 1062 (2001); Phys. Rev. B **65**, 045310 (2002).
- ⁶ B. I. Halperin, Jpn. J. Appl. Phys. Suppl. **26**, 1913 (1987).
- ⁷ G. Montambaux and M. Kohmoto, Phys. Rev. B **41**, 11417 (1990).
- ⁸ M. Kohmoto, B. I. Halperin, and Y. Wu, Phys. Rev. B **45**, 13488 (1992).
- ⁹ D. Pfannkuche and R. R. Gerhardts, Phys. Rev. B **46**, 12606 (1992).
- ¹⁰ A. Widom, Phys. Lett. **90A**, 474 (1982).
- ¹¹ P. Štředa, J. Phys. C **15**, L718 (1982).
- ¹² H. Aoki, Surf. Sci. **263**, 137 (1992).
- ¹³ M. Koshino, H. Aoki, and B. I. Halperin, Phys. Rev. B **66**, 081301(R) (2002).